

DEFORMATION QUANTIZATION IN COVARIANT HAMILTONIAN FIELD THEORY

G.SARDANASHVILY

Department of Theoretical Physics, Moscow State University
117234 Moscow, Russia
E-mail: sard@grav.phys.msu.su

Abstract

The deformation star product of smooth functions on the momentum phase space of covariant (polysymplectic) Hamiltonian field theory is introduced.

It is well known that, applied to field theory, the familiar symplectic technique of mechanics takes the form of instantaneous Hamiltonian formalism on an infinite-dimensional phase space. The finite-dimensional covariant Hamiltonian approach to field theory is vigorously developed from the seventies in its multisymplectic and polysymplectic variants, related to the two different Legendre morphisms in the first order calculus of variations on fibre bundles (see [1-5] for a survey).

Recall that, given a fibre bundle $Y \rightarrow X$ coordinated by (x^λ, y^i) , a first order Lagrangian L is defined as a semibasic density

$$L = \mathcal{L}\omega : J^1Y \rightarrow \bigwedge^n T^*X, \quad \omega = dx^1 \wedge \cdots \wedge dx^n, \quad n = \dim X,$$

on the jet manifold J^1Y of Y . Every Lagrangian L yields the Legendre map of J^1Y to the Legendre bundle

$$\Pi = \bigwedge^n T^*X \otimes_Y V^*Y \otimes_Y TX,$$

coordinated by $(x^\lambda, y^i, p_i^\lambda)$. It is provided with the canonical polysymplectic form

$$\Omega_Y = dp_i^\lambda \wedge dy^i \wedge \omega \otimes \partial_\lambda,$$

and is regarded as the polysymplectic momentum phase space of fields.

The multisymplectic momentum phase space of fields is the homogeneous Legendre bundle

$$Z_Y = T^*Y \wedge \binom{n-1}{T^*X},$$

coordinated by $(x^\lambda, y^i, p_i^\lambda, p)$ and equipped with the canonical multisymplectic form

$$d\Xi_Y = dp \wedge \omega + dp_i^\lambda \wedge dy^i \wedge \omega_\lambda.$$

The relationship between polysymplectic and multisymplectic phase spaces is given by the exact sequence

$$0 \rightarrow \Pi \times_X \binom{n}{T^*X} \hookrightarrow Z_Y \rightarrow \Pi \rightarrow 0,$$

where $Z_Y \rightarrow \Pi$ is a one-dimensional affine bundle. Given a section h of $Z_Y \rightarrow \Pi$, the pull-back $h^*\Xi_Y$ is a polysymplectic Hamiltonian form on Π .

It seems natural to generalize a Poisson bracket in symplectic mechanics to polysymplectic or multisymplectic manifolds in order to obtain the covariant canonical quantization of field theory. Different variants of such a bracket have been suggested (see the recent works [6,7]). The main difficulty is that the bracket must be globally defined, i.e., maintained under the bundle coordinate transformations.

Let us assume that a fibre bundle $Y \rightarrow X$ is affine. All realistic field models are of these type, and only vector and affine fields are quantized. One can introduce the bilinear differential operator

$$\Delta_\mu : (f, f') \mapsto f \overleftrightarrow{\Delta}_\mu f' = \partial_\mu f \partial_i f' - \partial_i f \partial_\mu f'$$

on smooth functions $f, g \in C^\infty(\Pi)$ on Π [8]. Let X be provided with a nowhere vanishing vector field ϑ . This condition implies the existence of a pseudo-Riemannian metric g on X such that integral curves of ϑ are time lines with respect to g . Let us consider the operator

$$\begin{aligned} \Delta : C^\infty(\Pi) \times C^\infty(\Pi) &\rightarrow C^\infty(\Pi), \\ \Delta : (f, f') &\mapsto f \overleftrightarrow{\Delta} f' = f [\overleftrightarrow{\Delta}_\mu \vartheta^\mu] f'. \end{aligned} \tag{1}$$

If $X = \mathbf{R}$, then $\vartheta = \partial_t$ and Δ (1) is the canonical Poisson bracket in Hamiltonian time-dependent mechanics [9,10].

Therefore, by analogy with the well-known Moyal product on symplectic manifolds, one can introduce the associative star product

$$f * f' = f \exp\left\{-\frac{i\hbar}{2} \overleftrightarrow{\Delta}\right\} f'$$

and the bracket

$$[f, f'] = f * f' - f' * f$$

of smooth functions on Π . This bracket provides deformation quantization of the momentum phase space of classical fields Π . Similarly to the familiar quantization in instantaneous Hamiltonian field theory, this deformation quantization is tangential to the integral curves of the vector field ϑ , i.e., to the time lines.

References

1. M.Gotay, A multisymplectic framework for classical field theory and the calculus of variations. I. Covariant Hamiltonian formalism. In: *Mechanics, Analysis and Geometry: 200 Years after Lagrange* (North-Holland, Amsterdam, 1991). 203-235.
2. G.Sardanashvily, *Generalized Hamiltonian Formalism for Field Theory. Constraint Systems* (World Scientific, Singapore, 1995).
3. G.Giachetta, L.Mangiarotti, and G.Sardanashvily, *New Lagrangian and Hamiltonian Methods in Field Theory* (World Scientific, Singapore, 1997).
4. M.Gotay, J.Isenberg and J.Marsden, E-print arXiv: physics/9801019.
5. G.Giachetta, L.Mangiarotti and G.Sardanashvily, Covariant Hamilton equations for field theory, *J. Phys. A* **32**, 6629 (1999); E-print arXiv: hep-th/9904062.
6. I.Kanatchikov, E-print arXiv: hep-th/0112263.
7. M.Forget, C.Paufler and H.Roemer, E-print arXiv: math-ph/0202043.
8. G.Sardanashvily, E-print arXiv: hep-th/9903220.
9. L.Mangiarotti and G.Sardanashvily, *Gauge Mechanics* (World Scientific, Singapore, 1998).
10. G.Sardanashvily, Hamiltonian time-dependent mechanics, *J. Math. Phys.* **39**, 2714 (1998)